

**THE UNITED REPUBLIC OF TANZANIA**  
**NATIONAL EXAMINATIONS COUNCIL**  
**ADVANCED CERTIFICATE OF SECONDARY EDUCATION EXAMINATION**

132/2

**CHEMISTRY 2**

**Time: 3 Hours**

**ANSWERS**

**Mwaka: 2003**

**Instructions**

1. This paper consists of a total of six questions
2. Answer five questions.

maktaba.tetea.org




1 (a)(i) Define the term dynamic equilibrium

Dynamic equilibrium is the state in a reversible chemical reaction in which the rate of the forward reaction is equal to the rate of the reverse reaction, and the concentrations of reactants and products remain constant over time, although the reactions continue to occur.

(ii) State the equilibrium law

The equilibrium law states that at constant temperature, the ratio of the product of the concentrations of the products raised to the power of their respective stoichiometric coefficients to the product of the concentrations of the reactants raised to the power of their stoichiometric coefficients is a constant known as the equilibrium constant ( $K_c$ ).

1 (b) Given the following system at equilibrium:



Predict the change of the concentration of  $\text{SO}_3$  if:

(i) The pressure of the system is increased

When pressure increases, the equilibrium will shift toward the side with fewer moles of gas to reduce the pressure. Since the forward reaction has fewer gas moles (3 moles  $\rightarrow$  2 moles), equilibrium shifts forward and concentration of  $\text{SO}_3$  increases.

(ii) A noble gas is added such that the pressure of the system increases but no volume changes occur

Adding a noble gas at constant volume does not change the partial pressures of the reacting gases, so there is no effect on the equilibrium and the concentration of  $\text{SO}_3$  remains unchanged.

(iii) More  $\text{SO}_3$  is added to the system

Adding more  $\text{SO}_3$  increases the concentration of product, and according to Le Chatelier's principle, the equilibrium will shift backward (reverse reaction favored) to reduce the concentration of  $\text{SO}_3$ .

(iv) The temperature of the system is increased

The forward reaction is exothermic (releases heat). Increasing temperature shifts equilibrium in the endothermic direction (backward), so the concentration of  $\text{SO}_3$  decreases.

1 (c) Consider the decomposition of phosphorus pentachloride:



Derive an expression which relates  $K_p$  and  $K_c$  for the above equilibrium where  $K_p$  and  $K_c$  are the equilibrium constants in terms of pressure and concentration respectively.

$$K_p = K_c \times (RT)^{\Delta n}$$


where  $\Delta n$  = moles of gaseous products - moles of gaseous reactants

In this case,  $\Delta n = 2 - 1 = 1$

So,  $K_p = K_c \times RT$

1 (d) At 500°C, the reaction between nitrogen and hydrogen to form ammonia has  $K_c = 8.0 \times 10^{-2}$ . Calculate its  $K_p$  value.

The reaction is:



$$\Delta n = 2 - (1+3) = -2$$

$$K_p = K_c \times (RT)^{\Delta n}$$

$$K_p = K_c \times (RT)^{-2}$$

Assuming  $R = 0.0821 \text{ atm}\cdot\text{dm}^3\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$  and  $T = 773 \text{ K}$  ( $500^\circ\text{C} = 773 \text{ K}$ )

$$K_p = 8.0 \times 10^{-2} \times (0.0821 \times 773)^{-2}$$

$$= 8.0 \times 10^{-2} \times (63.44)^{-2}$$

$$= 8.0 \times 10^{-2} \div 4023.4$$

$$= 1.99 \times 10^{-5}$$


$$K_p \approx 2.0 \times 10^{-5}$$

2 (a)

State Hess's Law of constant heat summation.

Hess's Law states that the total enthalpy change of a chemical reaction is the same, regardless of the number of steps in which the reaction occurs, provided the initial and final conditions are the same.

2 (b) By using Hess's Law, calculate the standard enthalpy of formation of methane ( $\text{CH}_4$ ) given that its standard enthalpy of combustion is  $-890 \text{ kJ mol}^{-1}$ , the standard enthalpy of combustion of carbon graphite is  $-393.5 \text{ kJ mol}^{-1}$ , and the enthalpy of formation of water is  $-285.9 \text{ kJ mol}^{-1}$ .



$$\Delta H_f(\text{CH}_4) = [\Delta H_c(\text{C}) + 2\Delta H_c(\text{H}_2)] - \Delta H_c(\text{CH}_4)$$

$$= (-393.5 + -571.8) - (-890)$$

$$= -965.3 + 890$$

$$= -75.3 \text{ kJ mol}^{-1}$$

2 (c) When 1.0 g of anhydrous copper (II) sulphate ( $\text{CuSO}_4$ ) was dissolved in a large amount of water, 0.418 kJ of heat were liberated. When 5.0 g of copper (II) sulphate pentahydrate crystals were dissolved in a large amount of water, 0.230 kJ of heat were absorbed. From this data calculate the heat change  $Z$  of the following reaction:



$$\text{Molar mass of CuSO}_4 = 159.5 \text{ g/mol}$$

$$\text{Moles of CuSO}_4 = 1.0 \div 159.5 = 0.00627 \text{ mol}$$

Heat per mole =  $0.418 \div 0.00627 = 66.67 \text{ kJ/mol}$

Molar mass of  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O} = 249.5 \text{ g/mol}$

Moles =  $5.0 \div 249.5 = 0.020 \text{ mol}$


Heat per mole =  $0.230 \div 0.020 = 11.5 \text{ kJ/mol}$

$\Delta H = -66.67 - (+11.5) = -78.17 \text{ kJ/mol}$

3 (a)

(i) Hard water becomes soft when washing soda ( $\text{Na}_2\text{CO}_3 \cdot 10\text{H}_2\text{O}$ ) is added


Because  $\text{Na}_2\text{CO}_3$  reacts with  $\text{Ca}^{2+}$  or  $\text{Mg}^{2+}$  ions forming insoluble  $\text{CaCO}_3$  or  $\text{MgCO}_3$  precipitates.



(ii) Iodine is sparingly soluble in water but dissolves readily in presence of iodide ions


Because  $\text{I}_2$  forms  $\text{I}_3^-$  ion with  $\text{I}^-$ :  $\text{I}_2 + \text{I}^- \rightarrow \text{I}_3^-$

(iii) Lead(IV) oxide forms bright yellow liquid in excess cold concentrated HCl



(iv) Effervescence of a colorless gas when  $\text{NaHCO}_3$  is added to copper(II) sulphate solution

$\text{NaHCO}_3$  reacts with acid producing  $\text{CO}_2$ :




(v) Mercury(I) iodide is soluble in potassium iodide solution but not in water


Due to formation of soluble complex:  $\text{Hg}_2\text{I}_2 + 2\text{I}^- \rightarrow 2[\text{HgI}_2]^-$

(vi) The pink solution of cobalt(II) chloride turns blue when concentrated HCl is added

Due to formation of blue  $\text{CoCl}_4^{2-}$  complex:




(vii) When water is sprayed into dry mixture of  $\text{SO}_2$  and  $\text{H}_2\text{S}$ , yellow solid is formed



(viii) Aqueous  $\text{AlCl}_3$  is acidic to litmus


$\text{Al}^{3+}$  hydrolyzes water forming  $\text{H}_3\text{O}^+$ :



(ix) HCl prevents precipitation of copper(II) sulphide from copper(II) salt solutions

Because HCl increases  $\text{H}^+$  concentration which suppresses  $\text{S}^{2-}$  formation.

(x) Effervescence occurs when ammonium chloride is added to warm sodium nitrite



4 (a)



(i) Electrochemical series: Arrangement of elements according to standard electrode potentials.

(ii) Electrochemical equivalent: Mass of substance deposited by unit electric charge.


(iii) Redox series: Arrangement of substances based on ease of oxidation or reduction.

(iv) Redox reaction: Chemical reaction involving transfer of electrons.

4 (b) Given:



(i) Feasible reaction: Zn reduces  $\text{Ni}^{2+}$  because Zn has lower  $E^\circ$  value.



(iii) Cell diagram:  $\text{Zn(s)} \mid \text{Zn}^{2+} \parallel \text{Ni}^{2+} \mid \text{Ni(s)}$

$$\text{(iv) } E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}} = -0.25 - (-0.76) = +0.51 \text{ V}$$

4 (c) Sodium chlorate(I) is converted by heat to sodium chlorate(V) and sodium chloride:



Oxidation: Cl from +1 to +5 ( $\text{NaClO}_3$ )

Reduction: Cl from +1 to -1 ( $\text{NaCl}$ )

5 (a) Give the meaning of volumetric analysis

Volumetric analysis is a quantitative method to determine concentration of solutions using titration based on volume measurements.

5 (b) Differentiate between:

(i) Molar solution = contains 1 mole of solute per litre

Normal solution = contains 1 equivalent per litre

(ii) Standard solution = known concentration



Primary standard = substance of high purity used to prepare standard solution

5 (c) Oxidation number of Cl:

(i) In  $\text{ClO}^- \rightarrow \text{Cl} = +1$

(ii) In  $\text{ClO}_3^- \rightarrow \text{Cl} = +5$

5 (d) Titration reaction:



$$\text{Moles of } \text{K}_2\text{Cr}_2\text{O}_7 = 1.85 \div 294 = 0.00629 \text{ mol}$$

$$\text{Concentration} = 0.00629 \div 0.25 = 0.0252 \text{ mol/L}$$

$$\text{Moles thiosulphate} = 0.0175 \times x$$

Using stoichiometry: 1 mol  $\text{Cr}_2\text{O}_7^{2-}$  requires 6 mol  $\text{S}_2\text{O}_3^{2-}$

0.00629 mol  $\text{Cr}_2\text{O}_7^{2-}$  needs 0.03774 mol thiosulphate  
 $x = 0.03774 \div 0.0175 = 2.156 \text{ mol/L}$

Electrons accepted = 6 per mole  $\text{Cr}_2\text{O}_7^{2-} \times 0.00629 = 0.03774 \text{ mol}$

6 (a)

- (i) Al is more metallic than B due to lower ionization energy and larger atomic radius.
- (ii) HI cannot be prepared from KI and  $\text{H}_2\text{SO}_4$  because  $\text{H}_2\text{SO}_4$  oxidizes  $\text{I}^-$  to  $\text{I}_2$ .
- (iii) Simple salts of  $\text{Cu}^+$  disproportionate to Cu and  $\text{Cu}^{2+}$  in aqueous solution.
- (iv) Carbon forms more stable hydrides due to small size and effective orbital overlap.
- (v) Li compounds have partial covalent character due to high polarizing power of  $\text{Li}^+$  ion.

6 (b) Group I elements have similar properties:


- React vigorously with water
- Form colorless solutions and hydroxides
- Show +1 oxidation state
- Have low ionization energy

7 (a)(i) What do you understand by the following terms?

Heat of reaction: It is the amount of heat evolved or absorbed during a chemical reaction under standard conditions.

(ii) Thermochemical equations: These are balanced chemical equations that include the enthalpy change ( $\Delta H$ ) of the reaction, showing whether the process is exothermic or endothermic.

7 (b) Given the following reactions:



(i) What do the symbols  $\Delta H_1^\circ$  to  $\Delta H_f^\circ$  above refer to?

$\Delta H_1^\circ$ : Enthalpy of atomization of Cd

$\Delta H_2^\circ$ : Enthalpy of ionization of Cd

$\Delta H_3^\circ$ : Enthalpy of sublimation of  $\text{I}_2$

$\Delta H_4^\circ$ : Bond dissociation enthalpy of  $\text{I}_2$

$\Delta H_5^\circ$ : Electron affinity of iodine

$\Delta H_f^\circ$ : Enthalpy of formation of  $\text{CdI}_2(\text{s})$

(ii) What is the sign for the enthalpy change corresponding to  $\Delta H_5^\circ$  above?

It is negative (–), since electron affinity is exothermic.

(iii) Define the standard enthalpy change corresponding to the following symbols:

$\Delta H_1^\circ$ : Enthalpy required to convert 1 mole of atoms in solid state to gaseous atoms.

$\Delta H_2^\circ$ : Enthalpy required to remove electrons from 1 mole of gaseous atoms.

$\Delta H_3^\circ$ : Enthalpy change when 1 mole of electrons is added to 1 mole of gaseous atoms.

$\Delta H_f^\circ$ : Enthalpy change when 1 mole of a compound is formed from its elements in standard states.

(iv) Calculate the value of  $\Delta H_f^\circ$  given:

$$\Delta H_1^\circ = +113 \text{ kJ/mol}$$

$$\Delta H_2^\circ = +2490 \text{ kJ/mol}$$

$$\Delta H_3^\circ = +199 \text{ kJ/mol}$$

$$\Delta H_4^\circ = +151 \text{ kJ/mol}$$

$$\Delta H_5^\circ = -314 \text{ kJ/mol}$$

$$\Delta H_l \text{ (lattice enthalpy)} = -2014 \text{ kJ/mol}$$

$$\Delta H_f^\circ = \Delta H_1^\circ + \Delta H_2^\circ + \Delta H_3^\circ + \Delta H_4^\circ + 2\Delta H_5^\circ + \Delta H_l$$

$$= 113 + 2490 + 199 + 151 + (2 \times -314) + (-2014)$$

$$= 113 + 2490 + 199 + 151 - 628 - 2014$$

$$= 3113 - 2642 = 471 \text{ kJ/mol}$$

8 (a) The members of the following pairs of isomeric compounds have different melting/boiling points. Indicate which member has the higher value and suggest reasons:

(i) Compound with  $-\text{OH}$  and  $-\text{NO}_2$  in para-position has higher melting point than ortho-isomer due to better packing in the lattice structure and symmetry.

(ii)

$\text{CH}_3\text{--C}(\text{OH})(\text{CH}_3)\text{--CH}_3$  has higher boiling point than  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$  due to stronger hydrogen bonding and higher branching leading to less surface area contact.

(iii)

$\text{CH}_3\text{CH}_2\text{OH}$  has higher boiling point than  $\text{CH}_3\text{OCH}_3$  due to hydrogen bonding in alcohols compared to weaker dipole interactions in ethers.

8 (b) From mass spectrum:

Molecular mass = 58

Empirical formula =  $\text{C}_3\text{H}_6\text{O}$

(i) Molecular formula =  $\text{C}_3\text{H}_6\text{O}$

(ii) Two possible structures:

Propanal ( $\text{CH}_3\text{CH}_2\text{CHO}$ ), Propanone ( $\text{CH}_3\text{COCH}_3$ )

8 (c) Liquid X tests:

| TEST | REAGENT                    | OBSERVATION                        |
|------|----------------------------|------------------------------------|
| A    | Sodium metal               | No reaction                        |
| B    | Bromine water              | No reaction                        |
| C    | 2,4-Dinitrophenylhydrazine | Orange precipitate (positive test) |
| D    | Tollen's reagent           | No reaction                        |

(i) Interpretation:

- Test A: no acidic H  $\rightarrow$  not alcohol
- Test B: no unsaturation  $\rightarrow$  not alkene
- Test C: carbonyl group present  $\rightarrow$  aldehyde or ketone
- Test D: negative  $\rightarrow$  not aldehyde  $\rightarrow$  hence ketone

(ii) Structure of X = Propanone ( $\text{CH}_3\text{COCH}_3$ )

9 (a) Explain briefly how the following conversions can be affected:

(i) A  $\rightarrow$   $\text{CH}_3\text{CH}_2\text{CH}_3$

Alkylation reaction with suitable reagent like  $\text{CH}_3\text{CH}_2\text{Cl} + \text{AlCl}_3$

(ii)

Aniline  $\rightarrow$  Bromoaniline

Electrophilic substitution using  $\text{Br}_2$

(iii)  $(\text{CH}_3)_3\text{N} \rightarrow [(\text{CH}_3)_3\text{N}]^+\text{OH}^-$

Reaction with  $\text{CH}_3\text{I}$  followed by aqueous silver oxide

9 (b) Distinguish between the following by chemical means:

(i) Phenol and Benzyl alcohol

Phenol reacts with  $\text{FeCl}_3$  to give violet complex; alcohol does not.

(ii)  $\text{CHCl}_3$  and  $\text{CH}_3\text{CCl}_3$

$\text{CHCl}_3$  gives isocyanide odor on reaction with  $\text{NaOH}$  and acetone;  $\text{CH}_3\text{CCl}_3$  does not.

(iii)  $\text{CH}_3\text{COCH}_3$  and  $\text{CH}_3\text{CH}_2\text{CHO}$

$\text{CH}_3\text{CH}_2\text{CHO}$  gives silver mirror with Tollen's reagent;  $\text{CH}_3\text{COCH}_3$  does not.

9 (c) Explain the following:

(i) Phenol is more acidic than ethanol due to resonance stabilization of phenoxide ion.


(ii) Trichloroethanoic acid is stronger than ethanoic acid due to inductive effect of Cl atoms.

(iii) Aniline is a weaker base than ethylamine due to delocalization of lone pair of nitrogen over the benzene ring.

10 (a) Give the products when ethyl alcohol reacts with concentrated sulphuric acid under the following conditions:


(i) 180°C

At 180°C, ethyl alcohol undergoes dehydration to form ethene:




(ii) 140°C

At 140°C, two molecules of ethyl alcohol condense to form diethyl ether:



10 (b) The following is part of a protein chain:



Draw the structure of two amino acids obtained on hydrolysis of this protein:

Amino acids on hydrolysis:

1.  $\text{CH}_3\text{-CH(NH}_2\text{)-COOH} \rightarrow$  Alanine

2.  $\text{CH}_3\text{CH}_2\text{-CH(NH}_2\text{)-COOH} \rightarrow$  Valine

10 (c)

1. Combustion data:

$\text{CO}_2$  from 1.173 g

Moles of C =  $1.173 \div 44 = 0.02666$  mol

mass of C =  $0.02666 \times 12 = 0.3199$  g

$\text{H}_2\text{O}$  from 0.240 g

Moles of H =  $0.240 \div 18 = 0.01333$  mol

mass of H =  $0.01333 \times 2 = 0.02666$  g

Mass of O =  $1.1 - (0.3199 + 0.02666) = 0.7534$  g

Moles of O =  $0.7534 \div 16 = 0.0471$  mol

Divide by smallest:

C =  $0.02666 \div 0.02666 = 1$

H =  $0.01333 \div 0.02666 = 0.5$

multiply by 2  $\rightarrow 1$

O =  $0.0471 \div 0.02666 \approx 1.77 \approx 3.5$

multiply by 2  $\rightarrow 7$

Empirical formula =  $C_2H_2O_7$   
empirical mass  $\approx 106$

2. Freezing point depression:

$$\Delta T = K_f \times m$$

$$0.186 = 1.86 \times (1.125 \div M) \times (1000 \div 125)$$

$$0.186 = 14.88 \div M$$

$$M = 14.88 \div 0.186 \approx 80 \text{ g/mol}$$

Approximate molecular mass = 80

Thus, molecular formula =  $C_2H_4O_3$  (most probable adjustment from empirical formula)