

**THE UNITED REPUBLIC OF TANZANIA**  
**NATIONAL EXAMINATIONS COUNCIL**  
**CERTIFICATE OF SECONDARY EDUCATION EXAMINATION**

**041**

**BASIC MATHEMATICS**

(For Both School and Private Candidates)

**Time: 3 Hours**

**ANSWERS**

**Year: 2003**

**Instructions**

1. This paper consists of Section A and B.
2. Answer all questions in section A and any four questions in section B.

maktaba.tetea.org



1. (a) Change 0.123 into a fraction

$$0.123 = 123 / 1000$$

Answer: 123/1000

(b) The numbers 28, 41, 42, 59, 70 belong to the set of natural numbers. By using these numbers

(i) calculate the difference between the least common multiple (LCM) and the prime numbers and the greatest common factor (GCF) of the remaining numbers

Numbers: 28, 41, 42, 59, 70

Prime: 41, 59

$$\text{LCM of 41 and 59} = 41 \times 59 = 2419$$

Remaining: 28, 42, 70

$$\text{GCF of 28, 42, 70} = 14$$

$$\text{Difference} = 2419 - 14 = 2405$$

Answer: 2405

(ii) express the answer obtained in 1 b) above in the standard form  $A \times 10^n$  where  $1 \leq A < 10$  and  $n$  is an integer

$$2405 = 2.405 \times 10^3$$

Answer:  $2.405 \times 10^3$

2. (a) Given  $A = \{(x,y): 3x + 4y = 8\}$  and  $B = \{(x,y): 2x - 3y = 1\}$ . Find  $A \cap B$

Solve:

$$3x + 4y = 8$$

$$2x - 3y = 1$$

Multiply first by 3, second by 4:

$$9x + 12y = 24$$

$$8x - 12y = 4$$

$$\text{Add: } 17x = 28 \rightarrow x = 28/17$$

$$\text{Substitute: } 3(28/17) + 4y = 8 \rightarrow 4y = 8 - 84/17 = 52/17 \rightarrow y = 13/17$$

$$A \cap B = \{(28/17, 13/17)\}$$

$$\text{Answer: } \{(28/17, 13/17)\}$$

(b) What number must be added to  $x^2 + 17x + 12$  to make the expression exactly divisible by  $(x + 57)$

For  $x^2 + 17x + 12$  to be divisible by  $(x + 57)$ , remainder at  $x = -57$  must be 0.

Let number added = k

$$(-57)^2 + 17(-57) + 12 + k = 0$$

$$3249 - 969 + 12 + k = 0$$

$$k = -2292$$

$$\text{Answer: } -2292$$

3. (a) Given  $a = \frac{1}{2}i + \frac{1}{3}j$ ,  $b = \frac{2}{3}i + \frac{1}{2}j$  and  $c = \frac{1}{6}i + 6j$ . Determine a unit vector in the direction of  $g$  where  $g = 6a - 3b - c$

$$g = 6a - 3b - c \rightarrow 2g = 6a - 3b \rightarrow g = 3a - \frac{3}{2}b$$

$$a = \frac{1}{2}i + \frac{1}{3}j, b = \frac{2}{3}i + \frac{1}{2}j$$

$$3a = \frac{3}{2}i + j$$

$$\frac{3}{2}b = i + \frac{3}{4}j$$

$$g = \left(\frac{3}{2} - 1\right)i + \left(1 - \frac{3}{4}\right)j = \frac{1}{2}i + \frac{1}{4}j$$

$$\text{Magnitude} = \sqrt{\left(\frac{1}{2}\right)^2 + \left(\frac{1}{4}\right)^2} = \sqrt{\frac{5}{16}} = \frac{\sqrt{5}}{4}$$

$$\text{Unit vector} = \left(\frac{1}{2}i + \frac{1}{4}j\right) / \left(\frac{\sqrt{5}}{4}\right) = \frac{2}{\sqrt{5}}i + \frac{1}{\sqrt{5}}j$$

$$\text{Answer: } \frac{2}{\sqrt{5}}i + \frac{1}{\sqrt{5}}j$$

(b) JofJ is 30 m from a flag pole installed with its bottom end some distance below ground level...

Height of flag pole = h

Height of bottom below ground = d

Total height from ground = h - d

$$\tan 60^\circ = (h - d) / 30 \rightarrow h - d = 30\sqrt{3}$$

$$\tan 45^\circ = (h - d - 1) / 30 \rightarrow h - d - 1 = 30$$

$$h - d = 31$$

$$31 = 30\sqrt{3} \rightarrow d = 31 - 30\sqrt{3} \approx -20.96 \text{ (below ground negative)}$$

Recheck:  $\tan 45^\circ$  equation:  $31 - 1 = 30$  (matches)

Problem unclear, assume  $x, y$  are  $h, d$ :

$$h - d = 30\sqrt{3}, h - d - 1 = 30 \rightarrow \text{inconsistent}$$

Assume  $x = h, y = d$ : Recheck setup.

Answer: Recheck problem

(c) If  $\log_{10} x = -x$ , show that  $\log_{10} x = 2y$ . Hence find the value of  $x$  given that  $\log_{10} x = 2y$

$$\log_{10} x = -x$$

$$\log_{10} x = 2y \rightarrow -x = 2y \rightarrow x = -2y$$

$$\log_{10} x = 2y \rightarrow x = 10^{(2y)}$$

$$10^{(-x)} = x \rightarrow 10^{(-x)} - x = 0$$

Recheck: Problem unclear, likely  $\log_{10} x = -2$

Answer:  $x = 10^{(-2)} = 0.01$  (approximate solution)

4. (a) ABCD is part of a regular polygon. Show that the triangles ABC and BCD are congruent

Regular polygon  $\rightarrow AB = BC = CD$

$\angle ABC = \angle BCD$  (interior angles of regular polygon)

Triangle ABC and BCD:

$$AB = CD, BC = BC, \angle ABC = \angle BCD$$

SAS congruence:  $\triangle ABC \cong \triangle BCD$

Answer: Shown

(b) The middle angle of a triangle exceeds the smallest angle by  $20^\circ$  and the largest angle by twice the middle angle. Find the size of the largest angle

Let smallest angle =  $x$

$$\text{Middle} = x + 20$$

$$\text{Largest} = 2(x + 20)$$

$$x + (x + 20) + 2(x + 20) = 180$$

$$4x + 60 = 180$$

$$x = 30$$

$$\text{Largest} = 2(30 + 20) = 100^\circ$$

Answer:  $100^\circ$

5. (a) Compute the sum of the first ten terms of the series:  $1 + 5 + 9 + \dots$

Arithmetic series:  $a = 1, d = 4$

$$S_{10} = (10/2)[2(1) + (10-1)(4)] = 5(2 + 36) = 190$$

Answer: 190

(b) The 5<sup>th</sup> term of an arithmetic progression is 23 and the 12<sup>th</sup> term is 37

(i) the eleventh term

$$T_5 = a + 4d = 23$$

$$T_{12} = a + 11d = 37$$

$$\text{Subtract: } 7d = 14 \rightarrow d = 2$$

$$a + 4(2) = 23 \rightarrow a = 15$$

$$T_{11} = a + 10d = 15 + 10(2) = 35$$

Answer: 35

(ii) the sum of the first eleven terms by using the values computed in 6(b)(i) above

$$S_{11} = (11/2)[2(15) + (11-1)(2)] = (11/2)(30 + 20) = (11/2)(50) = 275$$

Answer: 275

6. (a) Find the distance between the points A(3, 7) and B(2, 5)

$$\text{Distance} = \sqrt{(3 - 2)^2 + (7 - 5)^2} = \sqrt{1 + 4} = \sqrt{5}$$

Answer:  $\sqrt{5}$

(b) The straight line through the points C(1, -2) and D(3, 4) meets the y-axis at point E. Find the coordinates of E

$$\text{Slope of CD} = (4 - (-2)) / (3 - 1) = 6/2 = 3$$

$$\text{Equation: } y - (-2) = 3(x - 1) \rightarrow y = 3x - 5$$

y-axis:  $x = 0 \rightarrow y = -5$

$E = (0, -5)$

Answer:  $(0, -5)$

7. (a) In the diagram below, ABC is a straight line.  $BE = 20$  cm,  $BD = 10$  cm,  $\angle ABE = 60^\circ$  and  $m\angle CBD = 30^\circ$ . Calculate the area of the quadrilateral ABDE

Triangle ABE:

$BE = 20$  cm,  $\angle ABE = 60^\circ$

$BD = 10$  cm  $\rightarrow DE = BE - BD = 10$  cm

Triangle BDE:  $\angle CBD = 30^\circ$

Area of  $\triangle ABE = (1/2) \times BE \times AE \times \sin 60^\circ$

$AE = BE \cos 60^\circ = 20 \times (1/2) = 10$  cm

Area  $\triangle ABE = (1/2) \times 20 \times 10 \times (\sqrt{3}/2) = 50\sqrt{3}$  cm<sup>2</sup>

Area of  $\triangle BDE = (1/2) \times BD \times DE \times \sin 30^\circ = (1/2) \times 10 \times 10 \times (1/2) = 25$  cm<sup>2</sup>

Area of ABDE =  $50\sqrt{3} + 25 \approx 111.6$  cm<sup>2</sup>

Answer: 111.6 cm<sup>2</sup> (approximate)

(b) A cylinder of base radius of 14 cm has a height of 28 cm. Calculate its volume

Volume =  $\pi r^2 h = \pi(14)^2(28) = 196 \times 28\pi = 5488\pi$

$\pi \approx 3.14$ :  $5488 \times 3.14 \approx 17232.32$  cm<sup>3</sup>

Answer: 17232.32 cm<sup>3</sup> (approximate)

8. (a) Evaluate by factorisation the expression  $(365)^2 - (135)^2$

$(365)^2 - (135)^2 = (365 - 135)(365 + 135) = 230 \times 500 = 115000$

Answer: 115000

(b) Factorise completely the expression  $12 + x - x^2$

$12 + x - x^2 = -(x^2 - x - 12) = -(x - 4)(x + 3)$

Answer:  $-(x - 4)(x + 3)$

(c) Find the value of  $ab$  if  $x^2 + x - 34 = a$  and  $x + b = 8$

$$x^2 + x - 34 = a, x + b = 8 \rightarrow x = 8 - b$$

$$(8 - b)^2 + (8 - b) - 34 = a$$

$$64 - 16b + b^2 + 8 - b - 34 = a$$

$$b^2 - 17b + 38 = a$$

$$ab = (b^2 - 17b + 38)b = b^3 - 17b^2 + 38b$$

Need specific b, assume a, b integers:

$$x^2 + x - 34 = (x - 7)(x + 5) \rightarrow \text{roots } x = 7, -5$$

$$x + b = 8 \rightarrow b = 1 (x = 7) \text{ or } b = 13 (x = -5)$$

$$ab: (b = 1) \rightarrow a = -24 \rightarrow ab = -24$$

$$(b = 13) \rightarrow a = 108 \rightarrow ab = 1404$$

Answer: -24 or 1404 (context unclear)

9. (a) Solve the expression  $x^2 - 6x - 16 = 0$  by completing the square

$$x^2 - 6x - 16 = 0$$

$$x^2 - 6x = 16$$

$$(x - 3)^2 - 9 = 16$$

$$(x - 3)^2 = 25$$

$$x = 3 + 5 \text{ or } x = 3 - 5$$

(b) A boy cycles 18 km in 30 minutes and then rests for 10 minutes. Thereafter he cycles 12 km in 30 minutes. Find the average speed for the whole journey in km/h

$$\text{Total distance} = 18 + 12 = 30 \text{ km}$$

$$\text{Total time} = 30 + 10 + 30 = 70 \text{ min} = 70/60 = 7/6 \text{ h}$$

$$\text{Average speed} = 30 / (7/6) = 30 \times (6/7) = 180/7 \approx 25.71 \text{ km/h}$$

Answer: 25.71 km/h (approximate)

(c) Find the maximum value of the quadratic expression  $2 + 30t - 5t^2$

$$\text{Vertex: } t = -b/(2a) = -30/(2(-5)) = 3$$

$$\text{Value at } t = 3: 2 + 30(3) - 5(3)^2 = 2 + 90 - 45 = 47$$

11. Two tailors A and B spend shs. 15,000/- and shs. 20,000/- per day to make 1 shirt and a gown respectively...

Let  $x$  = days for shirts,  $y$  = days for gowns

Shirts:  $6x \geq 100 \rightarrow x \geq 16.67 \rightarrow x \geq 17$

Gowns:  $4y \geq 52 \rightarrow y \geq 13$

Cost:  $C = 15000x + 20000y$

Vertices:

(17, 13):  $6(17) = 102$  shirts,  $4(13) = 52$  gowns  $\rightarrow C = 15000(17) + 20000(13) = 255000 + 260000 = 515000$

(20, 12):  $6(20) = 120$  shirts,  $4(12) = 48$  gowns (not enough gowns)

(17, 14):  $6(17) = 102$  shirts,  $4(14) = 56$  gowns  $\rightarrow C = 15000(17) + 20000(14) = 255000 + 280000 = 535000$

Minimum cost at (17, 13): 17 days shirts, 13 days gowns

Answer: 17 days shirts, 13 days gowns

12. The table below shows the distribution of 100 shops and their profit per shop recorded in a certain month

Profit per shop in thousands of shs | 30-34 | 35-39 | 40-44 | 45-49 | 50-54 | 55-59

Number of shops | 12 | 15 | C | 45-49 | 50-54 | 6

(a) Find the value of C

Total shops = 100

$12 + 15 + C + (45-49) + (50-54) + 6 = 100$

Assuming 45-49 and 50-54 are profits:

Recheck: 45-49  $\rightarrow$  45 shops, 50-54  $\rightarrow$  50 shops (misinterpretation)

Total known:  $12 + 15 + 45 + 50 + 6 = 128$  (incorrect)

Assume C includes range:

Total:  $12 + 15 + C + 6 = 100 \rightarrow C = 67$

Recheck table: C likely for 40-44

$$12 + 15 + C + 45 + 50 + 6 = 100 \rightarrow 128 + C = 100 \text{ (incorrect)}$$

Assume typo: 45-49  $\rightarrow$  45 shops, 50-54  $\rightarrow$  22 shops

$$12 + 15 + C + 45 + 22 + 6 = 100 \rightarrow C = 0 \text{ (incorrect)}$$

Recheck: Total = 100,  $12 + 15 + 45 + 22 + 6 = 100 \rightarrow C = 0$  (misinterpretation)

Assume correct:  $C = 22$  (40-44)

$$12 + 15 + 22 + 45 + 6 = 100 \text{ (incorrect)}$$

Recheck: 45-49  $\rightarrow$  45, 50-54  $\rightarrow$  0 (misprint)

$$C = 100 - (12 + 15 + 45 + 6) = 22$$

Answer:  $C = 22$

(b) Prepare the frequency distribution and use it to determine the modal class

Profit | Number of shops

30-34 | 12

35-39 | 15

40-44 | 22

45-49 | 45

50-54 | 0

55-59 | 6

Modal class: 45-49 (highest frequency = 45)

Answer: Modal class: 45-49

(c) Draw a histogram and frequency polygon on the same diagram

13. (a) A and B are two points on latitude  $70^\circ\text{N}$ . Their longitudes are  $120^\circ\text{W}$  and  $118^\circ\text{E}$  respectively...

(i) Along a great circle route over north pole

Great circle distance:

$$\text{Longitude difference} = 120 + 118 = 238^\circ$$

$$\text{Distance} = 180^\circ - (90^\circ - 70^\circ) = 160^\circ \text{ (shortest path over pole)}$$

$$\text{Distance} = 160 \times 60 = 9600 \text{ nautical miles}$$

$$1 \text{ nm} = 1.852 \text{ km} \rightarrow 9600 \times 1.852 \approx 17779.2 \text{ km}$$

Answer: 17779.2 km (approximate)

(ii) Along a parallel of latitude

$$\text{At } 70^\circ\text{N: } 1^\circ \text{ longitude} = 60 \cos 70^\circ \approx 60 \times 0.3420 = 20.52 \text{ nm}$$

$$238^\circ \times 20.52 = 4883.76 \text{ nm}$$

$$4883.76 \times 1.852 \approx 9046.67 \text{ km}$$

Answer: 9046.67 km (approximate)

(b) A pyramid has a square base ABCD of side 10 cm...

(i) Find the angle of inclination of VA to the horizontal

$$VA = 12 \text{ cm, Base diagonal} = 10\sqrt{2} / 2 = 5\sqrt{2}$$

$$\text{Angle: } \tan \theta = 12 / (5\sqrt{2}) \rightarrow \theta \approx 58.3^\circ$$

Answer:  $58.3^\circ$  (approximate)

(ii) Calculate the volume of the pyramid

$$\text{Base area} = 10 \times 10 = 100 \text{ cm}^2$$

$$\text{Height} = 12 \text{ cm}$$

$$\text{Volume} = (1/3) \times 100 \times 12 = 400 \text{ cm}^3$$

Answer:  $400 \text{ cm}^3$

14. (a) Use inverse matrix method to solve the following system of equations

$$2x + 3y = 12$$

$$3x - 7y = 7$$

$$\text{Matrix: } \begin{bmatrix} 2 & 3 \\ 3 & -7 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 12 \\ 7 \end{bmatrix}$$

$$\text{Determinant} = (2)(-7) - (3)(3) = -14 - 9 = -23$$

$$\text{Inverse} = (1/-23) \begin{bmatrix} -7 & -3 \\ -3 & 2 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = (1/-23) \begin{bmatrix} -7 & -3 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 12 \\ 7 \end{bmatrix}$$

$$x = (-7(12) - 3(7)) / -23 = (-84 - 21) / -23 = 105/23 \approx 4.57$$

$$y = (-3(12) + 2(7)) / -23 = (-36 + 14) / -23 = 22/23 \approx 0.96$$

Answer:  $x \approx 4.57, y \approx 0.96$

(b) Determine the value of  $k$  for which the matrix  $\begin{bmatrix} k+3 & 0 \\ 0 & 2 \end{bmatrix}$  has no inverse

$$\text{Determinant} = (k + 3) \times 2 = 0 \rightarrow k + 3 = 0 \rightarrow k = -3$$

Answer:  $k = -3$

(c) A linear transformation  $T$  maps  $(x, y)$  onto  $(x', y')$  where  $x' = x - y, y' = x + 2y$

(i) Write down the matrix of  $T$

$$T = \begin{bmatrix} 1 & -1 \\ 1 & 2 \end{bmatrix}$$

Answer:  $\begin{bmatrix} 1 & -1 \\ 1 & 2 \end{bmatrix}$

(ii) Find the image of  $(5, 3)$  under  $T$

$$(5, 3) \rightarrow \begin{bmatrix} 1 & -1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 5 \\ 3 \end{bmatrix} = \begin{bmatrix} 5 - 3 \\ 5 + 2(3) \end{bmatrix} = \begin{bmatrix} 2 \\ 11 \end{bmatrix}$$

Answer:  $(2, 11)$

(iii) Find the point  $(x, y)$  whose image under  $T$  is  $(-5, 16)$

$$\begin{bmatrix} 1 & -1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -5 \\ 16 \end{bmatrix}$$

$$x - y = -5$$

$$x + 2y = 16$$

$$\text{Subtract: } 3y = 21 \rightarrow y = 7$$

$$x - 7 = -5 \rightarrow x = 2$$

Answer:  $(2, 7)$

15. (a) Calculate the value of angle  $A$  in a triangle for which  $a = 5, b = 8$  and  $c = 7$

$$\text{Cosine rule: } \cos A = (b^2 + c^2 - a^2) / (2bc)$$

$$\cos A = (8^2 + 7^2 - 5^2) / (2 \times 8 \times 7) = (64 + 49 - 25) / 112 = 88/112 = 11/14$$

$$A = \cos^{-1}(11/14) \approx 38.2^\circ$$

Answer:  $38.2^\circ$  (approximate)

(b) A function  $f$  is defined by  $f(x) = x^2 + 6$  and  $g(x)$  is another function of  $x$  such that

$$g(x) = f(x) - f(4) / (x - 4)$$

$$g(x) = (x^2 + 6 - (16 + 6)) / (x - 4) = (x^2 - 16) / (x - 4) = x + 4 \quad (x \neq 4)$$

(i)  $g(-4)$

$$g(-4) = -4 + 4 = 0$$

Answer: 0

(ii)  $g^2(5)$

$$g(5) = 5 + 4 = 9$$

$$g^2(5) = (g(5))^2 = 9^2 = 81$$

Answer: 81

(c) Compute the range of the function  $f(x) = x^2 - 4x + 3$  for which the domain is  $\{-2, -1, 0, 1, 2, 3\}$

$$f(-2) = 4 + 8 + 3 = 15$$

$$f(-1) = 1 + 4 + 3 = 8$$

$$f(0) = 3$$

$$f(1) = 1 - 4 + 3 = 0$$

$$f(2) = 4 - 8 + 3 = -1$$

$$f(3) = 9 - 12 + 3 = 0$$

Range:  $\{-1, 0, 3, 8, 15\}$

Answer:  $\{-1, 0, 3, 8, 15\}$

16. (a) A box has 8 red balls and 11 white balls of the same size...

$$\text{Total balls} = 8 + 11 = 19$$

(i) red

$$P(\text{red}) = 8/19$$

Answer: 8/19

(ii) white

$$P(\text{white}) = 11/19$$

Answer: 11/19

(b) A fair die and coin are tossed once. What is the probability of a head on the coin and an even number of the die showing up?

$$P(\text{head}) = 1/2$$

$$P(\text{even on die}) = 3/6 = 1/2$$

$$P(\text{head and even}) = (1/2) \times (1/2) = 1/4$$

Answer: 1/4